

TREATS UP TO 20,000 SQ.

Scotts, Accugreen and Edgeguard are registered trademarks of OMS Investments, Inc. Speedygreen is a registered trademark of VAS Inc. Speedygreen is a registered trademark of Everyis International B. V

Amdro Quick Kill, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

moo.lantaal@noisivibnabrag | 1707-262-008-f

Mos. DÄUMA

CENTRAL GARDEN & PET

header strips or stopped.

муви уои паув геаспед тре

feeding. Shut off spreader marks to ensure uniform **ACTIVE INGREDIENT**


*Bifenthrin 0.115% **OTHER INGREDIENTS** 99.885%

100.000% *Cis isomers 97% minimum, trans isomers 3% maximum. Contains 0.00115 lbs of Bifenthrin per pound. **KEEP OUT OF REACH OF CHILDREN** JTION See additional precautionary statements on back of bag

For outdoor use around residential sites only. For outdoor use only.









0.5 lb/500 sq. ft. is equivalent to 1 lb/1000 sq. ft. every 2 weeks only if there are signs of renewed insect activity. Evenly apply a 2-foot-wide band of granules next to the house/home foundation. Use 0.5 to 2.0 lbs. for each 500 sq. ft. (2 ft. x 250 ft.). If necessary, repeat applications **ТИЗИТАЗИТ ИЗВИВИТЕ**

HOME FOUNDATION

insect activity, but not more than every two weeks. Apply 0.5° to 1 lb. evenly over each 500 sq. ft. area. Repeat if there are signs of renewed This product can be applied just before planting or after plants emerge. **ORNAMENTALS & FLOWERS**

| In New York State, do make a single repeat applicati of renewed insect activity, but not sooner than two | on of this product if there are signs weeks after the first application. |
|---|---|
| In New York State, this product may NOT be applied equipment) to 300 feet (using aerial equipment) of o into coastal marshes. | bnuorg gnizu) təət OOT nirltiw nisıb tarlt emsərte ro zərlerısım latzacı |
| | the treated area after application. |

| Weevils (including Annual Bluegrass), Ants (including Argentine, Pharaoh (sugar), Imported Fire Ants, and Foraging Fire Ants) Millipedes Scorpions Ticks (including Brown Dog Tick) and ticks that may transmit Lyme Disease | uniformly apply 1 lb. per 250 sq ft. dayn or ld. lbs. per area of lawn or [4 lbs. per 1000, pp. lb]. For best results, thoroughly water the treated area after application. For Fire Ant Mounds: Uniformly applied to a ground or 3 teaspoons of granules to the surface of each mound. For best results apply in cool mound. For best results apply in cool weather, 65 to 80°F, in early weather, 65 to 80°F, in early mound. For lest results apply in cool weather, 65°F and 10°F, in early and 10°F |
|--|--|
| Southern Mole Cricket, Short-Winged Mole Cricket) | |
| Mites Mole Crickets (including Tawny Mole Cricket, | |
| Sround Pearl | |
| Ground Beetle | |
| Fleas (larvae and adults) | |
| Chiggers Crane Flies (larvae) | For best results in treating mole crickets, treat in late spring. |
| Annual Bluegrass Weevil (Hyperodes) | olog paiteost ai atlusos tand sod |
| Ants (including Carpenter, Red Harvester, Red Pavement, Odorous, Pyramid, Black Turf, White-footed, Crazy, Little Black, Small Honey, Lawn, Cornfield, Ghost, Thief, Acrobat, Big-Headed, and Allegheny mound) | Uniformly apply 1 to 2 lbs. per 500 sq. ft. or 2 to 4 lbs. per 1,000 sq. ft of lawn. For best results, thoroughly water the treated area after the application. |
| Wireworms | |

the lawn. Overlap wheel spreader the longest way of header strips. Operate the spreader. First apply two Mow lawn first if grass is tall enough to touch the bottom of Drop Spreader 6 **Z** %**S** Scotts AccuGreen overlapping occurs. previous one, so some about five feet from the Rotary солекаве таке еасh раss Scotts" Speedy Green" EdgeGuard" you stop. For complete **%9 %⊅ €** Shut off spreader anytime 11B 21B 41B Pounds of product per 1,000 sq.ft. DIRECTIONS **SPREADER** COVERS 5,000 SQ FT **NOITAJIJ99A** SPREADER SETTING

> **SPREADER SETTINGS** on the manufacturer's recommendations.

300534021 | AM1090_V3-6/23

Subscript used is first letter of run code on container.

EPA Est. No. 59144-GA-1_A, 92162-MO-001_B

1000 Parkwood Circle, Suite 700

Central Garden & Pet, Garden Division

responsibility for safety and use not in accordance with directions. Notice: To the extent consistent with applicable law, Buyer assumes all

disposal options. Never place unused product down any indoor or

If partly filled: Call your local solid waste agency for local

If empty: Place in trash or offer for recycling if available.

container. Do not reuse or refill this container. Do not pour or

PESTICIDE STORAGE: Store in a cool, dry, secure place. Do not

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable

dispose down-the-drain or sewer.

store near seeds, fertilizers, or foodstuffs.

STORAGE AND DISPOSAL

EPA Reg. No. 53883-194-90098

Atlanta, GA 30339

SPREADING INSTRUCTIONS
Follow Application Directions in the table below. Calibrate the spreader settings based

| ATZNI ƏNIQAƏRG | IOITOU | SN |
|--|--|---|
| broscoli, Brussels Sprouts, Chinese broccoli, Cabbage, Cauliflower, Cavalo broccoli, Chinese Cabbage, Kohlrabi, Chinese Mustard, Eggplant, Head Ettuce, Peppers (Bell and Non-Bell) | L | Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms |
| эАоґоіл | 2 | Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms |
| icurbit Vegetables ihayote, Citron melon, Cucumber, dible Gourd Muskmelon, Gantaloupe, asaba, Crenshaw melon, Golden melon, anta Claus melon, Snake melon) umpkin, Summer squash (Crookneck, callop, Straightneck, Zucchini) Winter quash (Butternut, Calabaza, Hubbard yatermelon | ε | Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms |
| ireen Peas, Edible Pods (Sugar Snap Peas, Snow Peas), Green Beans (Wax Ieans, Snap Beans), Dry Beans (Black Iyed Peas, Cow Pea) | ε | Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms |
| птоЭ 1ээм | L | Ants, armyworms, cutworms, crickets, earwigs, European corn borer, fall armyworm, grasshoppers, southwestern corn borer and weevils |
| səotsmo | ŀ | Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms |
| CROP | Days to wait after application till harvest | PESTS |

Apply a single application before planting or after plants emerge and work into the top 4 to 6 inches of soil. Use 1.0 lb of product for a 500 sq. ft. area. This product can be applied to the vegetables listed below. RESIDENTIAL VEGETABLE GARDEN

as rucy appears mornality water Webworms (including Sod Webworm, Grass Webworm, Tropical Sod Webworm) Two-lined Spittle Bugs Striped Grassworm Sowbugs Southwestern Corn Borer Springtails Spiders Silverfish Seed Maggots Pillbugs Mealybugs Lucerne Moth Leaf Hoppers Ladybeetles (including Convergent Ladybeetle, Seven-spotted Ladybeetle, Two-spotted Ladybeetle, Thirteen-spotted Ladybeetle, Twice-stabbed Ladybeetle, Asian Ladybeetle, Mealybug Ladybird) greenbug **Crasshoppers** Fruit Fly Fire Brat Fiery Skipper European Crane Fly Earwigs Dichondra Flea Beetle Cutworm, Granulate Cutworm, Variegated Cutworms (including Black Cutworm, Bronzed Spring Field Cricket, Northern Wood Cricket Western Striped Cricket, Eastern Striped Cricket, Sand Field Cricket, Vocal Field Cricket, Southern Southeastern Field Cricket, Texas Field Cricket, Cricket, Variable Field Cricket, Fall Field Cricket, Crickets (including House Cricket, Common Short-tailed Cricket, Arizona Cricket, Tropical sguð dənidə Chiggers Bristletail For best results, thoroughly water the treated area after application. Black Turfgrass Ataenius (adults) Billbugs (including Bluegrass Billbug, Hunting 1,000 sq. ft. area of lawn using a Armyworms (including Lawn Armyworm, Fall Armyworm) Uniformly apply 1 to 2 pounds over a PEST

APPLICATION RATE after use in measuring pesticides. NOTE: Food utensils such as tablespoons/teaspoons must not be used for food purposes Repeat application if there are signs of renewed insect activity but not more than once browned areas. For best results and to prevent further damage, treat the entire lawn, not just the after application. Do not use any other weed, disease, or insect control products for at least one week Water-in product after application. spreader settings. For best results, use a mechanical spreader. Spread this product uniformly over lawn at the labeled rates using the suggested

spots appear, or moths fly up from the grass when mowed).

SNWAJ This product is for use on lawns, around home foundations (perimeter treatment), ornamental and flower gardens, and vegetable gardens for the control of the insect pests **GENERAL INSTRUCTIONS**

Apply this product when insect activity or damage is first observed (lawn thins, brown

• For further information or to report suspected resistance contact your local sales representative. resistance-management and/or IPM recommendations for the specific site and pest problems in Contact your local extension specialist or certified crop advisors for any additional pesticide presence of resistance, consult with your local university specialist or certified pest control advisor. • Monitor after application for unexpected target pest survival. If the level of survival suggests the considers cultural, biological and other chemical control practices. Adopt an integrated pest management program for insecticide/acaricides use that includes scouting,
uses historical information related to pesticide use, crop rotation, record keeping, and which
uses historical information related to pesticide use, only for the period where both insecticides are active. unequal periods of residual insecticide activity may offer an insect resistance management benefit components have similar periods of residual insecticidal activity. Mixtures of insecticides with owt ent it setsestes are greated management benefits of an insecticide mixture are greatest if the two but they may still provide pest management benefits. Mixtures become less effective if resistance is already developing to one or both active ingredients, components for the targeted pest(s). When using mixtures, consider any known cross-resistance issues between the individual for insect resistance management. bahrammorer from strong the same IARI ambe of action classification are not recommended rates at which they are individually registered for use against thetarget species. Individual insecticides selected for use in mixtures should be highly effective and be applied at the by the Insecticide Resistance Action Committee (IRAC): pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same Use tank mixtures with insecticides/acaricides from a different group that are equally effective on among growing seasons, with different groups that control the same pests. • Rotate the use of this product or other Group 3A insecticides/acaricides within a growing season, or To delay insecticide/acaricide resistance, take the following steps: For resistance management, this product contains a Group A& insecticide/acaricides. Any insect population may contain individuals naturally resistant to this product and other Group A& insecticides/acaricides. The resistant individuals may dominate the insect population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management contacticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management contacticides/acaricides are used repeatedly in the same fields.

streams, marshes, ponds, estuaries, and commercial fish ponds). • Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, manner to limit dripping and avoid runoff onto untreated structural surfaces and plants. • To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a Treat surfaces to ensure thorough coverage but avoid runoff. • No not apply or irrigate to the point of runoff. except as a spot or crack-and-crevice treatment. • Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios • Do not allow the product to enter any drain during or after application. bodies, or aquatic habitat can occur. • Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies or anuatic pabitat can occur • Do not allow children or pets to enter the area until dry. \bullet Do not allow the product to enter any drain during or after application. • Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. • Do not apply when the wind speed is greater than 15 mph. Do not make applications during the rain.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DIKECTIONS FOR USE**

ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not **ENVIRONMENTAL HAZARDS**

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. **NOITUA** HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS

Note to Physician: This product is a pyrethroid. If large amounts have been ingested, milk, cream or other digestible fats and oils may increase absorption and so should be avoided. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information. • Call a poison control center or doctor for treatment advice. • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue lf in eyes Take off contaminated clothing.
 Rinze skin immediately with plenty of water for 15 – 20 minutes.
 Call a poison control center or doctor for treatment advice. lf on skin or clothing **GIA T2811**

moths fly up from the the grass and sbots appear, or ingredient into suggested spreader (ie. lawn thins, brown to release active over lawn using the activity or damage after application Spread uniformly Observe insect Water-in product STEP 1 STEP 2 **2LEb3 LAWN IN 3 EASY STEPS**

grass when mowed).

HOW TO TREAT YOUR GRANULES

